Data sources for FAO worldmaps of Koeppen climatologies and climatic net primary production

Jürgen Grieser, René Gommes, Stephen Cofield and Michele Bernardi

The Agromet Group, SDRN FAO of the UN, Viale delle Terme di Caracalla, 00100 Rome, Italy Contact: <u>Agromet@fao.org</u> or <u>juergen.grieser@rms.com</u>

August 2006

The <u>Koeppen climatologies</u> and the <u>climatic net primary production maps</u> of FAO are based on different periods and precipitation datasets. Here we provide the datasets in different formats. Furthermore some derived information like temperature of the coldest and warmest months, Martonnes aridity index and Gorczynskis continentality index are provided.

The original data are brought to a common grid based on <u>USGS gtopo30</u> and provided as tables in csv format (.5° resolution). For the users convenience the derived data are also provided as georeferenced data in <u>IDA/Windisp</u> format (5' resolution, resampled).

The table provides the links to the datasets used to derive the Koeppen climatologies and npp maps. Each of the files consists of 13 columns. The first column contains the gridpoint number, the remaining 12 columns contain the mean annual cycle of the variable at that grid point. In the case of temperature, it is the mean monthly temperature in °C or the standard deviation of temperature over the respective period. Precipitation is provided in mm per month. The meta data file consists of 4 columns with gridpoint number, longitude (in .01°), latitude (in .01°) and land fraction (in %).

All data are provided as comma separated value (csv) in $.5^{\circ}x.5^{\circ}$ resolution. The temporal standard deviation of the variable at the grid cell within the period is provided too. This allows a wide range of investigations. For example, it can be used to compare the average with the variability by estimating the coefficient of variability (standard deviation / average) in the case of precipitation. Furthermore it can be used to estimate uncertainty intervals for the average of each grid cell.

Comma Separated Value (csv)

	Full Period	Norm Period	Early Period	Late Period
	1951 – 2000	1961 – 1990	1951 – 1975	1976 - 2000
<u>CRU</u>	Average (2Mb)	Average	Average	Average
Temperature	<u>Standard</u>	Standard	<u>Standard</u>	Standard
	<u>Deviation</u>	Deviation	Deviation	Deviation
CRU	Average	Average	Average	Average
Precipitation	<u>Standard</u>	Standard	<u>Standard</u>	Standard
	Deviation	Deviation	Deviation	Deviation
GPCC Fulldata	Average	Average	Average	Average
Precipitation	<u>Standard</u>	Standard	<u>Standard</u>	<u>Standard</u>
	<u>Deviation</u>	Deviation	Deviation	<u>Deviation</u>
<u>GPCC</u>	Average	Average	Average	<u>Average</u>
VASClimO	Standard	Standard	Standard	Standard
Precipitation	Deviation	Deviation	Deviation	Deviation

The meta-data file with grid point coordinates is here.

For the annual mean temperature und the annual precipitation sum we also provide resampled georeferenced data in 5'x5' resolution as Windisp/IDA images.

Georeferenced annual Data					
	Full Period 1951 – 2000		Early Period 1951 – 1975		
CRU Temperature	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	
CRU Precipitation	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	
GPCC Fulldata Precipitation	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	
GPCC VASClimO Precipitation	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	

Download colour tables for IDA images of temperature, precipitation, number of months with temperature exceeding 10°C, and annual temperature amplitudes <u>here</u>.

Derived Temperature products					
	Full Period 1951 – 2000	Norm Period 1961 – 1990	Early Period 1951 – 1975	Late Period 1976 - 2000	
Mean monthly temperature of coldest month	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	
Mean monthly temperature of warmest month	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	
Mean annual temperature amplitude	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	
Number of months with temperature exceeding 10°C	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	<u>IDA</u>	
All as csv (700kb)	<u>CSV</u>	<u>csv</u>	<u>CSV</u>	<u>CSV</u>	

Aridity and Continentality


From the variety of existing indices to quantify aridity and continentality we only provide the aridity index of De Martonne (1926) and the continentality index of Gorczynski (1920).

Aridity indices provide a simple way to express the ratio of precipitation to evaporation. Since evaporation is rarely observed it is a common tradition to approximate it. In the approximation by De Martonne evaporation is set to mean annual temperature T_A in °C +10. The aridity index of De Martonne A_M is therefore defined as the ratio of the annual precipitation sum P_A in mm and the annual mean Temperature in °C +10. It is obvious that one disadvantage of this definition is that the equation has a pole at -10° C where the index is undefined. Lower temperatures lead automatically to negative indices. One may argue that the whole concept of aridity/humidity may not make much sense in cold regions. However, since we draw global maps we have to deal with this problem. In order to use the index world wide we define

$$A_M = \begin{cases} 100 & if T_A < -9.9^{\circ}C \\ P_A / (T_A + 10) & else \end{cases}$$

Note that the higher this coefficient is, the higher is the precipitation compared to evaporation and thus the less arid is the climate. This means that by definition a high aridity index means a humid climate while a low aridity index means an arid


climate. The following map shows the aridity index for the 50 year period from 1951 to 2000 based on temperature data of the CRU and precipitation data from GPCC VASCIimO. It can be downloaded as a bitmap <u>here</u>.



The continentality index of Gorczynski K_G is a simple but efficient way to estimate the influence of the ocean on the local climate. The index depends linearly on the annual temperature amplitude *A* (difference of monthly mean temperature of warmest and coldest month). However, *A* not only depends on the strength of the influence of the ocean but also on the annual cycle of incoming solar radiation. Since the amplitude of the annual cycle of incoming solar radiation depends on latitude, with a maximum in the polar regions, the inverse of the sine of the latitude φ gets in as well. The definition in the version of Gorczynski is

$$K_G = 1.7 \frac{A}{\sin \varphi} - 20.4$$

This original equation comes with some drawbacks. Since the sine approaches zero as the latitude approaches the equator, the values close to the equator tend to infinity. At the equator the definition breaks down. We therefore suggest not using the index values within a latitude range of plus/minus 10 degrees. In order to apply the definition also to the southern hemisphere we use the absolute of the latitude instead of the latitude itself. The following map shows the continentality index for the 50 year period from 1951 to 2000 based on temperature data of the CRU. It can be downloaded as a bitmap <u>here</u>.



Download colour tables for IDA images of De Martonne aridity index and of Gorczynski continentality index <u>here</u>.

De Martonne aridity index and Gorczynski continentality index					
Index	Full Period	Norm Period	Early Period	Late Period	
	1951 – 2000	1961 – 1990	1951 – 1975	1976 - 2000	
Gorczynski	<u>csv</u>	<u>csv</u>	<u>csv</u>	<u>csv</u>	
(CRU)	<u>IDA</u>	IDA	<u>IDA</u>	<u>IDA</u>	
De Martonne	<u>csv</u>	<u>csv</u>	<u>csv</u>	<u>csv</u>	
CRU	IDA	IDA	IDA	<u>IDA</u>	
De Martonne	<u>csv</u>	<u>csv</u>	<u>csv</u>	<u>csv</u>	
GPCC Fulldata	IDA	IDA	<u>IDA</u>	<u>IDA</u>	
De Martonne GPCC VASClimO	<u>csv</u> IDA	<u>csv</u> IDA	<u>csv</u> IDA	<u>csv</u> <u>IDA</u>	

Download this file as <u>pdf</u>.

References

Beck, C., J. Grieser and B. Rudolf, 2005: A New Monthly Precipitation Climatology for the Global Land Areas for the Period 1951 to 2000. *Klimastatusbericht 2004*, 181-190, DWD. [pdf]

De Martonne, E. (1941) : Nouvelle carte mondiale de l'indice s'aridité. Météorol. 1941, 3-26.

Gorczynski, W. (1920) : Sur le calcul du degré de continentalisme et son application dans la climatologie. Geogr. Annaler 2, 324-331.

Mitchell, T., and P. Jones, 2005: An improved method of constructing a database of monthly climate observations and associated high-resolution grids. Int. J. Climatol., 25, 693-712. <u>http://www.cru.uea.ac.uk/</u>

Rudolf, B., C. Beck, J. Grieser, U. Schneider, 2005: Global Precipitation Analysis Products of the GPCC. Internet publication at <u>http://gpcc.dwd.de/ [pdf]</u>